

[Nucleosides & Nucleotides, 11, 1731-1738 (1992)]

[Lab. of Medicinal Chemistry]

Synthesis of Thymidine Derivatives 3'-Modified with a Polar Three-atom Group as Potential Anti-HIV-1 Agents.

KOSAKU HIROTA*, HIROSHI HOSONO, YUKIO KITADE, YOSHIFUMI MAKI,

CHUNG K. CHU, RAYMOND H. SHINAZI, OSAMU MURAOKA

3'-Modified thymidine analogs, such as 3'-O-[(methylthio)thiocarbonyl]thymidine, 3'-O-thiocarbamoylthymidine, and *N*-(3'-deoxythymidin-3'-yl) phosphoramidates, were synthesized from thymidine and 3'-azido-3'-deoxythymidine (AZT) derivatives as potential anti-human immunodeficiency virus type 1 agents. No significant activity against HIV-1 was, however, observed with any of these compounds.

[J. Org. Chem., 57, 5268-5270 (1992)]

[Lab. of Medicinal Chemistry]

Convenient Method for the Synthesis of C-Alkylated Purine Nucleosides : Palladium-Catalyzed Cross-Coupling Reaction of Halogenopurine Nucleosides with Trialkylaluminums.

KOSAKU HIROTA*, YUKIO KITADE, YOSHITAKE KANBE, YOSHIFUMI MAKI

We have found that trialkylaluminums smoothly coupled with halogenopurine nucleosides. Thus, cross-coupling of 8-bromoadenosine protected with trimethylsilyl groups with trialkylaluminums in the presence of palladium catalyst, followed by deprotection, afforded the corresponding 8-alkyladenosines. This convenient method was applied to the synthesis of several C-alkylated purine nucleosides.

[J. Chem. Soc., Chem. Commun., 1992, 15-16]

[Lab. of harm. Synthetic Chemistry]

Ipsso Selectivity in the Reductive Iodonio-Claisen Rearrangement of Allenyl (p-methoxyaryl) iodinanenes.

MASAHITO OCHIAI*, TAKAO ITOH, YUKIO MASAKI

Allenyl (aryl) iodinanenes, generated from p-methoxy (diacetoxyiodo) arenes by the reaction with propyn-2-yl (trimethyl) silanes in the presence of BF₃·Et₂O in dichloromethane, undergo reductive ipso iodonio-Claisen rearrangement selectively at -20 °C yielding ipso-substituted propynylarenes.